THE UNITED METHODIST CHURCH MEDICAL SUMMARY REPORT OF MINISTERIAL CANDIDATE Form 103

Candidate's Name:	
To the Board of Ordained Ministry:	
Please indicate here, the name/address of the boa	rd officer who will receive this report.
Name:	
Address:	
	OR THE RELEASE OF TION – COMPLETED BY CANDIDATE
Candidate Name:	Birth Date:
(annual conference) Bo	(physician) to disclose to the pard of Ordained Ministry the following information with (candidate) for the purpose of evaluation by The United
	this consent at any time except to the extent that action vill expire sixty (60) days after the date treatment is
otherwise protected by federal as well as state lav	be disclosed from records whose confidentiality is v. Any of the above requested information may include diagnosis and treatment of psychological disorders, as
confidentiality is protected by federal law. Federa any further disclosure of it without the specific wi	mation has been disclosed to you from records whose all regulations (42 CFR Part 2) prohibit you from making ritten consent of the person to whom it pertains, or as ral authorization for the release of medical or other
Signature of Candidate	Date
Witness	Date
	♥ @gbhem URL gbhem.org f facebook.com/gbhem ADDRESS 1001 19th Avenue, South

Nashville, TN 37212



SUMMARY REPORT - COMPLETED BY PHYSICIAN

Comments for physician:

Candidate's Name:

Complete the summary report. The United Methodist Church assumes you are completing this information based on a current physical examination of the candidate. Screening guidelines are provided for reference as needed.

This person is a candidate for ministry in The United Methodist Church. Among other requirements, this includes being able to typically work a full-time week – with periodic weeks requiring longer work hours. Those serving in ministry will encounter situations that require the ability to cope with conflict and stress. Job-related tasks range from office work and traveling from site to site to communicating with and relating to a variety of people and managing multiple tasks simultaneously, among other responsibilities.

Date of Physical Exam:
Check One:
CHECK OHE.
Based on the physical exam I completed, this candidate appears to be healthy. I have no concerns about his/her physical fitness for ministry.
Based on the physical exam I completed, this candidate has some health concerns that are summaraized below.
Summary of Concerns:
@gbhem URL gbhem.org

Nashville, TN 37212

Typical treatment(s) for this condition could potentially include (medication, surgery, lifestyle medication, intervention by specialist, frequent monitoring, etc.):				
	_			
	_			
	_			
Questions to ask, or conve include:	rsation that a committee	might have, to address	s these concerns could	



Examining P	rovider:
Address:	
Phone:	
Fax:	
Signature:	Date:

STAMP



EXAMINATION STANDARDS*

As a part of the ministry application process, The United Methodist Church requires each candidate to "present a satisfactory certificate of good health" by a physician on the prescribed form. Disabilities are not to be construed as unfavorable health factors when a person with disability is capable of meeting the professional standards and is able to render effective service.... (*The Book of Discipline*, ¶¶ 315.6c, 324.8, 347.3, 357.7, 355.3, 368.5).

The following lists show standard screening practices to be considered in an assessment of physical health. Additionally, the physician may choose to make recommendations to the candidate as needed. While the candidate's physician should make the final determination regarding the need for specific medical tests as related to the overall health and needs of the candidate, The United Methodist Church seeks a summary report from the physician upon completion of a physical examination of the candidate that provides an assessment of the candidate's physical ability to perform the required work of ministry.

NOTE: DO NOT RECORD SCREENING RESULTS ON THIS FORM.

Screening

Height and weight (periodically)

Blood pressure

Alcohol and tobacco use

Depression (if appropriate follow-up is available)

Diabetes mellitus (patients with hypertension)

Dyslipidemia (total and HDL cholesterol): men ≥35 y; men or women ≥20 y who have

cardiovascular risk factors; measure every 5 y if normal

Colorectal cancer screening (men and women 50-75 y)

Mammogram every 1 to 2 y for all women \ge 40 y. Evaluation for BRCA testing in high-risk women only.

Papanicolaou test (at least every 3 y until age 65 y)

Chlamydial infection (sexually active women ≤25 y and older at-risk women)

Routine voluntary HIV screening (ages 13-64 y)

Bone mineral density test (women ≥65 y and at-risk women 60-64 y)

AAA screening (one time in men 65-75 y who have ever smoked)

Counseling—Substance Abuse

Tobacco cessation counseling

Alcohol misuse: brief behavioral counseling; alcohol abuse: referral for specialty treatment

Counseling—Diet and Exercise

Behavioral dietary counseling in patients with hyperlipidemia, risks for CHD and other diet-related chronic disease

Regular physical activity (at least 30 minutes per day most days of the week) Intensive counseling/behavioral interventions for obese patients

AAA = abdominal aortic aneurysm; BRCA = breast cancer susceptibility gene; CHD = coronary heart disease.

* Based on recommendations from the U.S. Preventive Services Task Force.

Key Points

- The U.S. Preventive Services Task Force recommends routine periodic screening for hypertension, obesity, dyslipidemia (men ≥35 years), osteoporosis (women ≥65 years), abdominal aortic aneurysm (one-time-screening), depression, and HIV infection.
- The U.S. Preventive Services Task Force recommends routine periodic screening for colorectal cancer (persons 50-74 years of age), breast cancer (women ≥40 years), and cervical cancer.
- The U.S. Preventive Services Task Force recommends that all pregnant women be screened for asymptomatic bacteriuria, iron-deficiency anemia, hepatitis B virus, and syphilis.
- The U.S. Preventive Services Task Force recommends against screening for hemochromatosis; carotid artery stenosis; coronary artery disease; herpes simplex virus; or testicular, ovarian, pancreatic, or bladder cancer.
- Outside of prenatal, preconception, and newborn care, genetic testing should not be performed in unselected populations because of lower clinical validity; potential for false positives; and potential for harm, including "genetic labeling."
- For patients for whom genetic testing may be appropriate, referral for genetic counseling should be provided before and after testing.
- A human papillomavirus vaccine series is indicated in females ages 9 through 26 years, regardless of sexual activity, for prevention of cervical cancer.
- A single dose of tetanus-diphtheria-acellular pertussis (Tdap) vaccine should be given to adults ages 19 through 64 years to replace the next tetanus-diphtheria toxoid (Td) booster.
- A zoster (shingles) vaccine is given to all patients 60 years and older regardless of history of prior shingles or varicella infection.
- Asymptomatic adults who plan to be physically active at the recommended levels do not need to consult with a physician prior to beginning exercise unless they have a specific medical question.
- Smoking status should be determined for all patients.
- Patients who want to quit smoking should be offered pharmacologic therapy in addition to counseling, including telephone quit lines.
- Routine screening is recommended to identify persons whose alcohol use puts them at risk.
- For management of alcohol abuse and dependence, referral for specialty treatment is recommended; for management of alcohol misuse, brief behavioral counseling may be useful.



- Clues for chemical dependency include unexpected behavioral changes, acute intoxication, frequent job changes, unexplained financial problems, family history of substance abuse, frequent problems with law enforcement agencies, having a partner with substance abuse, and medical sequelae of drug abuse.
- Condom use reduces transmission of HIV, Chlamydia, gonorrhea, Trichomonas, herpes virus, and human papillomavirus.
- It is important to ask about domestic violence when patients present with symptoms or behaviors that may be associated with abuse.
- When an abusive situation is identified, address immediate safety needs.